

NAME: \_\_\_\_\_ Score \_\_\_\_\_ /100

Please print

**SHOW ALL YOUR WORK IN A NEAT AND ORGANIZED FASHION**

Course Average \_\_\_\_\_

1. T F A matrix is a rectangular array of numbers.
2. T F Matrix addition is commutative.
3. T F Matrix multiplication is commutative.
4. T F The inner product of two matrices is a matrix.
5. T F If A and B are square matrices with the same dimension, then  $\det(AB) = \det(A)\det(B)$ .
6. T F The rule for the det function is a recursive rule.
7. T F If A is a square matrix and  $\det(A) = 0$ , then A does not have an inverse.
8. T F If two matrices have the same order, their product is defined.
9. T F Only square matrices can be multiplied.
10. T F Every square matrix has a determinant.
11. T F Division of matrices is defined as a multiplication.
12. If a matrix has m rows and n columns, the order of the matrix is \_\_\_\_\_.
13. If a matrix has only one row, it is called a \_\_\_\_\_ matrix.
14. If a matrix has only one column, it is called a \_\_\_\_\_ matrix.
15. Two matrices are equal if they have the same \_\_\_\_\_ and their corresponding \_\_\_\_\_ are equal.
16. The \_\_\_\_\_ matrix of order n is the matrix whose main diagonal entries are 1 and all other entries are 0.
17. The determinant is a function whose domain is the set of \_\_\_\_\_ matrices and whose range is the \_\_\_\_\_ numbers.
18. The matrix consisting of the coefficients (in the same order) of a system of equations is called the \_\_\_\_\_ matrix.
19. Write the  $3 \times 3$  identity matrix \_\_\_\_\_.
20. The entry in the 34 position of the matrix AB is the inner product of row \_\_\_\_\_ in matrix A and column \_\_\_\_\_ in matrix B.
21. Write the coefficient matrix of the following system of equations: 
$$\begin{cases} -2x + 2y - 4z = 1 \\ 2x - 5y - z = 6 \\ 4x + 2y - 3z = 5 \end{cases}$$
22. Write a matrix equation which is equivalent to the system 
$$\begin{cases} 3x + 7y - 4z = 5 \\ 2x - 3y - 5z = 3 \\ -2x + 3y - 6z = 0 \end{cases}$$

23. Complete the statements of the **Elementary Row Operations**:

a. Interchange two \_\_\_\_\_.

b. \_\_\_\_\_ a row by a non-zero \_\_\_\_\_ and \_\_\_\_\_ that row with the \_\_\_\_\_.

c. Add a \_\_\_\_\_ of a row to another \_\_\_\_\_ and replace \_\_\_\_\_ but not both of the \_\_\_\_\_ with that \_\_\_\_\_.

24. Calculate  $\det \begin{pmatrix} \begin{bmatrix} 5 & 1 \end{bmatrix} \\ \begin{bmatrix} 6 & 2 \end{bmatrix} \end{pmatrix} =$

25. Calculate the inner product  $\begin{bmatrix} 2 & -2 & 3 \end{bmatrix} \begin{bmatrix} -4 \\ -1 \\ 2 \end{bmatrix} =$

26. Perform the addition  $\begin{bmatrix} 2 & 5 & 2 \\ -1 & 0 & 6 \\ 4 & -2 & 1 \end{bmatrix} + \begin{bmatrix} -8 & 5 & -1 \\ -5 & 6 & -4 \\ 1 & 1 & 0 \end{bmatrix} =$

27. Calculate  $\det \begin{pmatrix} \begin{bmatrix} 3 & 6 & 4 \\ 1 & 0 & 2 \\ -1 & -6 & 5 \end{bmatrix} \end{pmatrix}$

28. The inverse of  $A = \begin{bmatrix} -2 & -3 & 1 \\ -3 & -3 & 1 \\ -2 & -4 & 1 \end{bmatrix}$  is the matrix  $A^{-1} = \begin{bmatrix} 1 & -1 & 0 \\ 1 & 0 & -1 \\ 6 & -2 & -3 \end{bmatrix}$

Use this information to solve the system  $\begin{cases} -2x - 3y + z = 2 \\ -3x - 3y + z = 0 \\ -2x - 4y + z = 3 \end{cases}$

29. Supply the missing entries by performing the indicated elementary row operation.

$$\begin{bmatrix} 2 & -2 & 1 & 0 \\ 3 & -3 & 0 & 1 \end{bmatrix} \xrightarrow{\frac{1}{3}R_2 \rightarrow R_2} \begin{bmatrix} \square & \square & \square & \square \\ \square & \square & \square & \square \end{bmatrix}$$

30. Supply the missing entries by performing the indicated elementary row operation

$$\begin{bmatrix} -2 & 3 & \frac{1}{5} & 0 \\ 6 & -3 & 0 & 1 \end{bmatrix} \xrightarrow{3R_1 + R_2 \rightarrow R_2} \begin{bmatrix} \square & \square & \square & \square \\ \square & \square & \square & \square \end{bmatrix}$$

31. Perform the multiplication:

$$\begin{bmatrix} 2 & 3 & -4 \\ -1 & -3 & 3 \end{bmatrix} \begin{bmatrix} 3 & -2 \\ 4 & -1 \\ -2 & 1 \end{bmatrix} =$$

32. You do not need to do any computations. Simply fill in the blanks to describe the process for finding the inverse of a matrix.

To find the inverse of the matrix  $A = \begin{bmatrix} 5 & 7 & 4 \\ 3 & -1 & 3 \\ 6 & 7 & 5 \end{bmatrix}$

Begin by adjoining the \_\_\_\_\_ matrix to obtain the matrix  $\begin{bmatrix} 5 & 7 & 4 & 1 & 0 & 0 \\ 3 & -1 & 3 & 0 & 1 & 0 \\ 6 & 7 & 5 & 0 & 0 & 1 \end{bmatrix}$  with order \_\_\_\_\_

The next step is to get a \_\_\_\_\_ in the \_\_\_\_\_ position.

Then use that \_\_\_\_\_ to get \_\_\_\_\_ everywhere else in the \_\_\_\_\_

At this point the matrix will have been converted to  $\begin{bmatrix} 1 & 7/5 & 4/5 & 1/5 & 0 & 0 \\ 0 & -26/5 & 3/5 & -3/5 & 1 & 0 \\ 0 & -7/5 & 1/5 & -6/5 & 0 & 1 \end{bmatrix}$

The next step is to get a \_\_\_\_\_ in the \_\_\_\_\_ position.

Then use that \_\_\_\_\_ to get \_\_\_\_\_ everywhere else in the \_\_\_\_\_

At this point the matrix will have been converted to  $\begin{bmatrix} 1 & 0 & 25/26 & 1/26 & 7/26 & 0 \\ 0 & 1 & -3/26 & 3/26 & -5/26 & 0 \\ 0 & 0 & 1/26 & -27/26 & -7/26 & 1 \end{bmatrix}$

The next step is to get a \_\_\_\_\_ in the \_\_\_\_\_ position.

Then use that \_\_\_\_\_ to get \_\_\_\_\_ everywhere else in the \_\_\_\_\_

At this point the matrix will have been converted to  $\begin{bmatrix} 1 & 0 & 0 & 26 & 7 & -25 \\ 0 & 1 & 0 & -3 & -1 & 3 \\ 0 & 0 & 1 & -27 & -7 & 26 \end{bmatrix}$

The inverse of A is matrix  $A^{-1} =$

Which has order \_\_\_\_\_